arXiv:cond-mat/0411550v1 [cond-mat.stat-mech] 22 Nov 2004

Glass glass transition and new dynamical singularity points in an analytically solvable p-spin glass like model

Antonio Caiazzo, Antonio Coniglio, Mario Nicodemi
Dipartimento di Scienze Fisiche, Universitá "Federico II",
INFM-Coherentia and INFN Napoli, via Cintia, I-80126 Napoli, Italy

We introduce and analytically study a generalized p-spin glass like model that captures some of the main features of attractive glasses, recently found by Mode Coupling investigations, such as a glass/glass transition line and dynamical singularity points characterized by a logarithmic time dependence of the relaxation. The model also displays features not predicted by the Mode Coupling scenario that could further describe the attractive glasses behavior, such as aging effects with new dynamical singularity points ruled by logarithmic laws or the presence of a glass spinodal line.

PACS numbers: 64.70.Pf, 05.50.+q 64.60.Ht

Recent Mode Coupling Theory (MCT) [1] investigations discovered [2] a new kind of glasses, the so called attractive glasses, characterized by the presence of a glass/glass transition line where higher order dynamical singularity points are located. As MCT calculations triggered intense experimental researches [3, 4], their limit is that, by definition, they can correctly describe only the region outside the glass. In this paper, we introduce a schematic p-spin glass like model exhibiting a glass/glass transition which we can analytically investigated to establish the proper scenario of the glassy phase. This leads to new testable predictions on the aging dynamics in that region, where new types of singularities are found.

In MCT attractive glasses the potential between the particles presents attraction for some range besides the hard core repulsion [2]. Its volume fraction - temperature phase diagram shows two branches of the glass transition line, one due to the repulsive part of the potential, extending to high temperatures for volume fractions near the close packing value, and one related to the attraction, extending to low volume fractions up to meet the liquid/gas spinodal. For sufficiently narrow widths of the potential well one of the two branches interestingly enters in the glass region giving rise to a glass/glass transition line, ending in a higher order dynamical singularity point. This special point in the MCT framework is characterized by a logarithmic time dependence of the relaxation. The theory applies to attractive colloids; actually, some evidence of the MCT results has been found by recent experiments [3, 4]. Nevertheless, for some aspects, the approach used in [2] turns out to be inadequate. It is known in fact that MCT is valid only for equilibrium dynamics and cannot be trivially extended inside the glassy region, where the glass/glass line itself is located and aging phenomena are present. Another problem arises when the phase coexistence interferes with the glass transition line. In this case MCT does not allow a self-consistent description of the structural relaxation and the gas/liquid spinodal line is usually obtained *a priori*, not taking into account the presence of the glassy phase. The purpose of the present work is to begin to cure these inconsistencies. Exploiting the strong analogies between glassy systems and the class of 'discontinuous' spin glasses noticed in the last years [5], we introduce a schematic solvable model providing a coherent picture of the glass/glass line, its off-equilibrium dynamics and singularity points.

The model consists of a combination of two (or more) p-spin models [6, 7, 8] diluted through density variables [9]. We analytically solve its Langevin dynamics and for comparison with MCT inside the glassy region we also use the Mode Coupling approximation. We show that: (A) The glass/glass transition line does not coincide with that predicted by MCT but follows a different curve (Figs. 1,2). (B) Along the glass/glass line (see Fig. 1) in the early regime the stationary autocorrelation function $\Phi(t-t')$ exhibits, like in MCT, a power law relaxation except at its endpoint A_3 where is logarithmic. (C) In the long time regime, where MCT predicts a plateau, we find an interesting aging dynamics. In particular, two more dynamical singularity points, B_3 and B_3' (one for each kind of glass, see Fig.1) characterizes the relaxation in the aging regime. On one side of the glass/glass line from C to B_3 (resp. B_3' on the other side, see inset in Fig. 1) the long times relaxation is a power law, except right at B_3 where it is logarithmic (analogously for B_3'). From B_3 (resp. B_3') to A_3, the aging behavior is spin glass-like [5]. (D) Finally, at the A_4 singularity (i.e., in the case where A_3 coincides with point C in Fig. 1) we show that the logarithmic decay of the correlation function in the stationary regime $\Phi\left(t-t'\right)$ is followed by an aging regime ruled again by a logarithmic law. (E) We calculate in a consistent way the density-temperature phase diagram and the spinodal lines (see Fig. 1) and find they are influenced by the presence of the glassy phase.

The Model. We consider a lattice gas model, where particles ($n_i = 0, 1$) interact in groups of p via their spins ($s_i = \pm 1$) through quenched random couplings $J_{i_1 \cdots i_p}$, with zero mean and variance $p! J_p^2 / \left(2N^{p-1}\right)$, and also in groups of r through (nonrandom) coupling constants K_r

(typically below we take two or three values of p and one value of r). The mean field Hamiltonian is given by:

$$H = -\sum_p \sum_{i_1<\cdots<i_p} J_{i_1\cdots i_p} s_{i_1} n_{i_1} \cdots s_{i_p} n_{i_p} - \sum_r \frac{r! K_r}{2N^{r-1}} \sum_{i_1<\cdots<i_r} n_{i_1} \cdots n_{i_r} - \mu \sum_i n_i \quad (1)$$

where μ is the chemical potential. The spherical version of the model (1) is here considered: we introduce two new spin variables s_{ai} $(a = 1, 2)$, defined by $s_i = s_{1i}$, $n_i = (s_{1i} s_{2i} + 1)/2$, and enforce for these ones spherical constraints ($\sum_i s_{ai}^2 = N$) [10]. To exclude the possibility of continuous transitions, not related to a glass transition, the restriction $p > 2$ is taken [6]. The two spin fields s_{ai} $(a = 1, 2)$ evolve via usual Langevin equations with thermal noises $\xi_{ai}(t)$, having zero mean and variance $\langle \xi_{ai}(t) \xi_{bj}(t') \rangle = 2T \delta_{ab} \delta_{ij} \delta(t - t')$. Due to its quadratic nature the model is exactly solvable. It has a gas, a liquid and a glassy phases, respectively called P^-, P^+ and G (see Fig. 1). We focus here on the dynamics, and in particular on the correlation function $C(t, t') = \sum_{ia} \langle s_{1i}(t) s_{ai}(t') \rangle / (2N)$, the related response function $G(t, t')$ and the density $d(t) = C(t, t)$. Standard functional techniques [7, 11] allow to derive the following dynamical equations for these quantities after a rapid quench at $t = 0$ from high temperature $(t \geq t')$:

$$\frac{\partial C(t,t')}{\partial t} = -\frac{z(t)}{2} C(t,t') + 2T G(t',t) + \frac{1}{2} \int_0^{t'} du\, \varphi'(C(t,u))\, G(t',u) + \frac{1}{2} \int_0^{t} du\, \varphi''(C(t,u))\, G(t,u)\, C(u,t') \quad (2)$$

$$\frac{\partial G(t,t')}{\partial t} = -\frac{z(t)}{2} G(t,t') + \frac{\delta(t-t')}{2} + \frac{1}{2} \int_{t'}^{t} du\, \varphi''(C(t,u))\, G(t,u)\, G(u,t') \quad (3)$$

$$d'(t) = (z(t) + 2\mu + 2\chi'(d(t)))(1 - d(t)) - T \quad (4)$$

Here we define $\varphi(x) = (1/2) \sum_p J_p^2 x^p$, $\chi(x) = (1/2) \sum_r K_r x^r$ and $z(t) = z_1(t) - \mu - \chi'(d(t))$ with $z_1(t) = z_2(t)$ the two time-dependent Lagrange multipliers related to the spherical constraints (the prime stands for the first derivative). The above equations completely define the dynamical model, given the initial density $d(0)$. In the following we consider the large times limit, where one-time quantities reach stationary values: $d = d(t \to \infty)$ and $z = z(t \to \infty)$.



FIG. 1: The model dynamical phase diagram in the density-temperature plane (the parameters are here: $p_1 = 3$, $p_2 = 11$, $J_3^2 = .74$, $J_{11}^2 = 4$ and $r = 12$, $K_{12} = -12$). One can have a fluid (P) phase consisting of gas (P^-) and liquid (P^+), or a glass (G) phase. Continuous lines depict the liquid/glass (P^+/G) and glass/glass (G/G) transition lines, the spinodal of the fluid (P^--sp and P^+-sp) and that of the glass (G-sp). The **inset** shows an enlargement of the region around the G/G transition line. The dotted line represents the incorrect G/G line predicted by MCT. On the G/G line (full line from C to A_3), along with its endpoint A_3, there are new dynamical singularity points, B_3, B_3', characterized by a logarithmic relaxation. Also the crossing point C between the two branches of the P^+/G transition line is shown.

MCT calculations. Assuming equilibrium dynamics, we have that $C = C(t-t')$ and $G = G(t-t')$ $\forall t, t'$ and the Fluctuation-Dissipation theorem (FDT) holds: $TG(\tau) = -C'(\tau)$ (with $\tau = t - t'$). Under these hypotheses, Eqs. (2,3) yield the long time schematic MCT equation [12] for the correlator $\Phi(\tau) = C(\tau)/d$,

$$\tau_0 \frac{d\Phi(\tau)}{d\tau} + \Phi(\tau) + \int_0^\tau d\tau' m(\tau - \tau') \frac{d\Phi(\tau')}{d\tau'} = 0 \quad (5)$$

where $m(\tau) = \varphi'(C(\tau))\, d/T^2$. In the standard MCT notation [1] the kernel $m(\tau)$ is usually written as $m(\tau) = F(\Phi(\tau), v)$ with $F(f, v) = \sum_n v_n f^n$; this allows to establish a mapping relating T, d, J_p and the usual MCT control parameters v_n, given by $v_n = p J_p^2 d^p / (2T^2)$ where $n = p - 1$. If $p_1, p_2, \ldots$ denote the values assumed by p, we deal with the so-called $F_{p_1-1, p_2-1, \ldots}$ MCT model. The non-ergodicity parameter $f = \Phi(\tau \to \infty)$ is obtained by solving the bifurcation equation $Z(f) = 1$ (i.e., the $\tau \to \infty$ limit of Eq. (5)), where

$$Z(f) = \frac{1}{1-f} - F(f, v) = \frac{1}{1-f} - \sum_p \frac{p J_p^2 d^p}{2T^2} f^{p-1} \quad (6)$$

The inequality $1 \leq Z(\Phi(\tau))$, deriving from the decreasing character of $\Phi(\tau)$, implies that f is the largest solution of the bifurcation equation [1]. The conditions $Z(f_c) = 1$ and $Z'(f_c) = 0$ determine the transition line in the plane $d - T$ (see Fig. 1). The P^+/G transition, where f jumps from zero to a nonzero value, f_c, corresponds to singularity points of type A_2 [1]. MCT Eq. (5) describes the relaxation when this critical line is approached from the P^+ phase: $\Phi(\tau)$ decays in two steps characterized by power laws with critical exponents determined by the parameter $\lambda = 1-(1-f_c)^3 Z''(f_c)/2 < 1$ [1]. Inside the glassy region, for a set of values of the parameters v_n, an other line of A_2 singularities appear (dotted line in the inset of Fig. 1) where two non zero solutions for the dynamical order parameter, f, are allowed. This line defines the glass-glass (G/G) transition, where a first glass, f_{c1}, transforms discontinuously into a second one, f_{c2} [1, 2, 13]. Its endpoint, where $\lambda = 1$, i.e., $Z''(f_c) = 0$, results to be an higher order singularity A_3 (with $f_{c1} = f_{c2}$). To have a G/G line in our model, at least two values of p are needed [14]: as an example, for $p_1 = 3$ the lowest other possible value is $p_2 = 11$ (the case of Fig.s 1,2).

The glassy region and the new G/G line. Eq.(5), i.e., MCT, is correct only out of the glassy phase (for $f = 0$), where the equilibrium dynamics assumption holds. Thus, the above results on the G/G line are not correctly described by this theory. Within our model, we discuss now the more complex non-equilibrium equations governing the glassy phase, the correct position of the G/G transition and the new aging behavior along the G/G line. In the glassy phase, as in the usual p-spin glass model, the relaxation of $\Phi(t,t')$ can be split in two parts [5, 8, 11]: in the first, describing the so called FDT regime ($t \simeq t'$), $\Phi(t,t')$ is a function, $\Phi_{FDT}(\tau)$, of the times difference $\tau = t - t'$; the second part for $t \gg t'$ corresponds to an aging regime described by a function $\Phi_{AG}(t,t')$. In the FDT regime, i.e., the approach of the correlator to the plateau f_c, $\Phi_{FDT}(\tau)$ satisfies the following equation:

$$\tau_0 \frac{d\Phi(\tau)}{d\tau} + Z\Phi(\tau) + (1-Z) + \int_0^\tau d\tau' m(\tau-\tau') \frac{d\Phi(\tau')}{d\tau'} = 0 \tag{7}$$

where the existence of a well defined aging solution fixes Z to the minimum value (not necessarily equal to 1) satisfying the stability requirement $Z(f) \le Z(\Phi_{FDT}(\tau))$, i.e., the one corresponding to the marginal stability condition $Z'(f) = 0$ [8, 11, 15] (see Fig. 2). Note that Eq.(7) coincides with (5) only for $Z = 1$. The G/G transition line results thus to be modified with respect to MCT and located by the conditions $Z(f_{c1}) = Z(f_{c2})$ and $Z'(f_{c1}) = Z'(f_{c2}) = 0$ (see Fig.s 1,2). Its endpoint A_3 (see Fig. 1), is characterized, like in MCT, by one more condition: $f_{c1} = f_{c2} = f_c$ with $Z''(f_c) = 0$, corresponding to $\lambda = 1$.

In the aging region, the correlator $C_{AG}(t,t') = \Phi_{AG}(t,t')\, d$ and the response function $G_{AG}(t,t')$ obey a generalized FDT relation, $TG_{AG}(t,t') = x\partial C_{AG}(t,t')/\partial t'$, where $x \le 1$ is a constant with the physical meaning of the ratio between the bath temperature and an effective temperature T_{eff}, describing the out of equilibrium configurations $x = T/T_{eff}$ [5].



FIG. 2: The function $Z(f)$ defined in Eq.(6) is plotted for several decreasing temperatures T (from the top to the bottom) and fixed density. For high T the liquid solution (P^+) is given by the value $f = 0$ corresponding to the lowest value of Z namely $Z = 1$. By lowering T the glass transition T_c is reached when the minimum of $Z(f)$ approaches the level $Z = 1$ (bold line marked P^+/G). For $T \le T_c$ the plateau f_c is given by the value of f where $Z(f)$ has the minimum. For low enough T, $Z(f)$ develops a second minimum. The G/G transition is reached when the two minima have the same depth (bold line marked G/G). Note that in MCT approximation this would be reached instead when the second minimum approaches the level $Z = 1$ (line marked G/G^{MCT}).

The dynamics along the G/G line. On the G/G line two glasses coexist corresponding to two different values of the plateau, f, and thus two different values of $x = x(f)$ and $\lambda = \lambda(f)$ (with $0 \le x \le 1$ and $0 \le \lambda \le 1$), except at the endpoint A_3 where the glasses coincide. As shown below, along the line the value of λ determines the properties of the dynamics in the FDT regime, namely the approach to the plateau, while the ratio λ/x determines the aging regime, i.e., the departure from the plateau. Four different behaviors are found along the G/G line (see Fig. 1): the case $\lambda/x < 1$, corresponding to the points from C to B_3 (resp. B_3', on the other side of the line, see below); the new dynamical singularity points B_3 and B_3' where $\lambda/x = 1$; the case $\lambda/x > 1$ extending from B_3 (resp. B_3') to the endpoint A_3; and the singular point A_3 itself where $\lambda/x > 1$, but $\lambda = 1$.

More precisely the approach to the plateau in the FDT regime, along the entire G/G line, is given by $\Phi_{FDT}(\tau) - f_c \sim \tau^{-\beta}$, where the exponent β is given by $\Gamma^2(1-\beta)/\Gamma(1-2\beta) = \lambda$ [5, 8, 11] (Γ is Euler gamma function). At the endpoint A_3 where $\lambda = 1$ this is replaced by a logarithmic behavior given by $\Phi_{FDT}(\tau) - f_c \sim 1/\ln^2 \tau$ [1, 2, 13].

The departure from the plateau in the aging regime, along the G/G line from the crossing point C up to the two new points B_3 and B_3', one for each side of the line, is given by a power law:

$$\Phi_{AG}\left(t'+\tau,t'\right)-f_c\sim-\left(\tau/\mathcal{T}\left(t'\right)\right)^{\alpha} \tag{8}$$

with $\Gamma^2\left(1+\alpha\right)/\Gamma\left(1+2\alpha\right)=\lambda/x$ [8]. Here the t'-dependence is contained in $\mathcal{T}\left(t'\right)\sim t'$ [16].

At the two new dynamical singular points, B_3 and B_3', where $\lambda/x=1$, this power law is replaced by a logarithmic behavior, one for each side of the G/G line. For instance, by approaching B_3 from the right (the side where the plateau value f_c is higher) one has

$$\Phi_{AG}\left(t'+\tau,t'\right)-f_c\sim 1/\ln^2\left(\tau/\mathcal{T}\left(t'\right)\right) \quad . \tag{9}$$

From the point B_3 to the endpoint A_3 (analogously from B_3' to A_3 on the other side of the G/G line), $\lambda/x>1$. This implies that the present one-step Replica Symmetry Breaking solution should not hold [11] and, instead, a spin glass-like aging dynamics should be found in such a region [5].

Interestingly the length of the G/G line can be varied and, in a version of the model with three p-values [14], the A_3 endpoint singularity can be made to coincide with the crossing point C (see Fig. 1). In this particular case the A_3 becomes an A_4 singularity, as also found by MCT [2] (the simplest model with such a singularity is found to be $p_1=3, p_2=4, p_3=11$ [14]). In this case the FDT regime is logarithmic $\Phi_{FDT}\left(\tau\right)-f_c\sim 1/\ln\tau$, as in MCT [1, 2, 13], as well as the aging regime:

$$\Phi_{AG}\left(t'+\tau,t'\right)-f_c\sim 1/\ln\left(\tau/\mathcal{T}\left(t'\right)\right). \tag{10}$$

Spinodal lines. We show now that the glass transition curve P^+/G intersects the fluid spinodal line determining the existence of a glass spinodal line. The two spinodals are obtained by enforcing the vanishing local stability conditions in Eq. (4):

$$\beta^2\varphi''\left(d\right)+2\beta\chi''\left(d\right)-\frac{1}{\left(1-d\right)^2}-\frac{1}{d^2\left(1-f\right)^2}=0 \tag{11}$$

The P^-/P^+-spinodal corresponds to a solution, $T(d)$, with $f=0$, as for the G-spinodal solution the value of f of the glassy phase is given by the marginal stability condition $Z'\left(f\right)=0$. The two lines, completing the dynamical phase diagram, together with their range of validity are shown in Fig. 1.

In conclusion, the model we have introduced is a simple extension of p-spin glass models [6, 7, 8] for the glass transition. Our model exhibits a glass/glass transition line which can be analytically investigated. Actually, it turns out to be different from the G/G line derived by equilibrium MCT and to be characterized by new dynamical singularity points and off-equilibrium properties. These results can help to shed deeper light on the properties of attractive glasses, such as colloidal suspensions.

Work supported by MIUR-PRIN 2002, FIRB 2002, CRdC-AMRA, INFM-PCI, EU MRTN-CT-2003-504712.

---

[1] W. Götze, in *Liquids, freezing and glass transition*, Hansen J.P., Levesque D., Zinn-Justin editors (North Holland, Amsterdam, 1991); W. Götze and M. Sperl, Phys. Rev. E 66 (2002) 11405; J. Phys. Condens. Matter 15(11) (2003) S869; M. Sperl, cond-mat 0310772

[2] K. Dawson *et al.*, Phys. Rev. E 63 (2001) 11401; E. Zaccarelli, G. Foffi, K. Dawson, F. Sciortino and P. Tartaglia, Phys. Rev. E 63 (2001) 31501;

[3] F. Mallamace *et al.*, Phys. Rev. Lett. 84 (2000) 5431; W.R. Chen, S.H. Chen and F. Mallamace, Science 300 (2003) 619

[4] T. Eckert and E. Bartsch, Phys. Rev. Lett. 89 (2002) 125701; K.N. Pham *et al.*, Science 296 (2002) 104; W.C.K. Poon, K.N. Pham, S.U. Egelhaaf and P.N. Pusey, J. Phys. Condens. Matter 16 (2003) S269

[5] J.P. Bouchaud, L.F. Cugliandolo, J. Kurchan and M. Mezard, 'Out of equilibrium dynamics in spin glasses and other glassy systems' in *Spin glasses and random fields*, edited by A.P. Young (World Scientific, Singapore, 1997)

[6] Th. M. Nieuwenhuizen, Phys. Rev. Lett. 74 (1995) 4289; S. Ciuchi and A. Crisanti, Europhys. Lett. 49 (2000) 754

[7] H. Sompolinsky and A. Zippelius, Phys. Rev. B 25 (1982) 6860; T.R. Kirkpatrick and D. Thirumalai, Phys. Rev. B 36 (1987) 5388

[8] L.F. Cugliandolo and J. Kurchan, Phys. Rev. Lett. 71 (1993) 173;

[9] J. Arenzon, M. Nicodemi and M. Sellitto, J. Phys. I France 6 (1996) 1143; *ibid.* 7 (1997) 945

[10] A. Caiazzo, A. Coniglio and M. Nicodemi, Phys. Rev. E 66 (2002) 46101; Europhys. Lett. 65(2) (2004) 256

[11] L.F. Cugliandolo and P. Le Doussal, Phys. Rev. E 53 (1996) 1525

[12] The full MCT equation also contains a second-order time derivative term [1]. Since it becomes negligible at large times, it is often ignored in literature, as explained in [5].

[13] W. Götze and M. Sperl, Phys. Rev. E 66 (2002) 11405; cond-mat 0403278. M. Sperl, cond-mat 0310772

[14] detailed calculations will be reported in a longer paper by A. Caiazzo, A. Coniglio and M. Nicodemi, in preparation

[15] H. Kinzelbach and H. Horner, J. Phys. I (France) 3 (1993) 1329; J. Phys. I (France) 3 (1993) 1901; S. Franz and M. Mezard, Physica A 209 (1994) 1; A. Crisanti, H. Horner and H.J. Sommers, Z. Phys. B 92 (1993) 257

[16] More precisely $\mathcal{T}\left(t'\right)=h\left(t'\right)/h'\left(t'\right)$,where $h\left(t\right)$ is a time reparametrization increasing function (for the p-spin glass model one can assume $h\left(t\right)\sim t$, see [8]).